



Transformation of uranium under Fe(III)-reducing conditions: Reduction of U(VI) by biogenic green rust

Edward O'Loughlin and Kenneth M. Kemner

Argonne National Laboratory

Michelle M. Scherer

University of Iowa

ERSP Annual PI Meeting

April 4, 2006



*A U.S. Department of Energy laboratory
managed by The University of Chicago*



Project Members

Molecular Environmental Science Group - ANL

Ed O'Loughlin, Ken Kemner, Shelly Kelly, Maxim Boyanov, and Bruce Ravel

Electron Microscopy Center - ANL

Russ Cook

University of Iowa

Michelle Scherer, Philip Larese Casanova, Sharon Smith, and Justine Harrison

Coe College

Martin St.Clair


Special Thanks To

John Zachara (PNNL), and Jim Fredrickson (PNNL), and Eric Roden (UW)




Abiotic, Microbial, and Microbially Mediated Reduction


Abiotic




Microbial



Microbially Mediated





Dissimilatory Fe(III) Reduction



✓ Effective reductants for many organic and inorganic contaminants


Fe(II)-Fe(III) Hydroxide (Green Rust)

- Layered Structure
- General Composition: $[Fe^{II}_{(6-x)}Fe^{III}_x(OH)_{12}]^{x+} [(A)_{x/n} \bullet yH_2O]^{x-}$
- Formed via numerous biotic and abiotic processes
- Plays a central role in the redox cycling of Fe in many aquatic and terrestrial environments
- Capable of reducing many organic and inorganic contaminants (e.g., halogenated hydrocarbons, Se(VI), Cr(VII), Hg(II), Tc(VI), U(VI), etc.)






Dissimilatory Fe(III) Reduction

Experimental System




160-mL serum vial
100 mL Minimal Medium
75 mM e^- donor
80 mM Fe(III)
 $\sim 5 \times 10^9$ cells mL^{-1}



Shewanella Surveyed for DIR of $g\text{-FeOOH}$


Shewanella

- **g-Proteobacteria**
- **One of the most phylogenetically diverse genus of *Bacteria***
- **Facultatively anaerobic**
- **Remarkable flexibility wrt range of e^- acceptors for anaerobic respiration**
- **Physiological diversity reflected in range of habitats (mesophilic, psychrotolerant, psychrophilic, barophilic)**
- **Isolated from a broad range of environments (marine and freshwater, sediments, terrestrial subsurface)**
- **Commonly found at oxic/anoxic boundaries**




0.10



Reduction of Lepidocrocite by *Shewanella* spp.



96 d






Lepidocrocite Reduction by CN32



Characterization of Biogenic Solids



S. putrefaciens CN32




Green Rust produced by all but *S. denitrificans*





e⁻ Donor Effects on DIR of g-FeOOH by CN32

Fe(II) production resulting from dissimilatory reduction of lepidocrocite by *S. putrefaciens* CN32




e- Donor Effect on Solid-Phase Fe(II) Products



U(VI) Interaction with Biogenic Green Rust


Experimental System



Biogenic GR
Pasteurized 70 °C
Washed/sonicated
58 mM Fe(II)
Add 500 µM U(VI)



pH
U speciation
U XAFS
Total U





Within 48 h > 98% of added U associated with solids
Adsorption or Reduction???????



XAFS (X-ray Absorption Fine Structure)




X-ray Absorption Fine Structure Spectroscopy



- Short-range structural probe – provides information on oxidation state as well as the number, type(s), and distances of neighboring atoms (typically 1-2 “shells”) along with an indication of structural disorder




U(VI) Interaction with Biogenic Green Rusts



Modeling of EXAFS Data

Model based on structure of uraninite (UO_2)



Neighboring atom	Microbe type	Number of neighbors	Distance to Neighbors (?)	Disorder in distance to neighbors ($\times 10^{-3} \text{ Å}^2$)
U-O1	CN32	6.8 ± 0.4	2.31 ± 0.01	13.9 ± 1.7
	ANA3	7.2 ± 0.5		
	Alga	6.8 ± 0.4		
U-O2	CN32	1.2 ± 0.4	2.67 ± 0.05	13.9 ± 1.7
	ANA3	0.8 ± 0.5		
	Alga	1.2 ± 0.4		
U-Fe	CN32	1.9 ± 0.9	3.41 ± 0.02	10.4 ± 5.0
	ANA3	1.8 ± 0.7		
	Alga	2.0 ± 0.8		
U-U	CN32	3.3 ± 1.3	3.77 ± 0.03	10.4 ± 5.0
	ANA3	4.8 ± 1.4		
	Alga	3.8 ± 1.1		
U-O3	CN32	6.6 ± 2.5	4.43 ± 0.03	20.8 ± 10.0
	ANA3	9.6 ± 2.8		
	Alga	7.7 ± 2.2		



U(IV) structure in U-BioGR is similar to nanoparticulate uraninite (UO_2)




Comparison of Shewanella species




Results similar with green rusts formed by different species





U(VI) Reduction by Green Rust



Particle size from U-U coordination



Modeling of EXAFS $\text{U-GR}_{\text{SO}_4}$
data predicted ~1-2 nm UO_2 particles



TEM imaging
revealed
nanoscale
 UO_2 particles
(~2-5 nm)

Summary

Lepidocrocite Bioreduction

- Variability in lepidocrocite bioreduction among *Shewanella* species
- Broad range of *Shewanella* species can form green rust
- Carbonate green rust dominant product with formate, H₂, lactate, and N-acetylglucosamine
- Siderite dominant product with serine and pyruvate

U(VI) reduction

- U(VI) reduced to U(IV) in presence of biogenic green rust
- Molecular clusters of uraninite-like U(IV) in close association with Fe
- Results similar for green rusts formed by different species
- Biogenic green rust may be a reactive phase for contaminant reduction in Fe(III)-reducing environments

For all this and MORE, see our poster TONIGHT!

